arXiv:math/0402393v2 [math.GT] 4 Mar 2005

STRONGLY-CYCLIC BRANCHED COVERINGS OF KNOTS VIA $(g,1)$-DECOMPOSITIONS

PAOLA CRISTOFORI, MICHELE MULAZZANI, AND ANDREI VESNIN

Abstract. Strongly-cyclic branched coverings of knots are studied by using their $(g,1)$-decompositions. Necessary and sufficient conditions for the existence and uniqueness of such coverings are obtained. It is also shown that their fundamental groups admit geometric g-words cyclic presentations.

1. Introduction

A $(g,1)$-decomposition of a knot in a closed orientable 3-manifold is a genus g Heegaard splitting of the manifold, such that the knot is trivially split in two arcs by the Heegard surface (a more detailed definition will be given in Section 2). Any knot K in a 3-manifold N admits a $(g,1)$-decomposition, for a certain $g \geq h(N)$, where $h(N)$ is the Heegaard genus of N. In the following a knot admitting a $(g,1)$-decomposition will be called a $(g,1)$-knot. The particular case of $(1,1)$-knots has recently been investigated in several papers from different points of view (see references in [3]).

In this paper we study strongly-cyclic branched coverings of knots, by using topological and algebraic properties of their $(g,1)$-decompositions. Some of the obtained results naturally generalize results from [2] and [7] concerning $(1,1)$-knots.

In Section 2, we prove a representation theorem of $(g,1)$-knots in terms of Heegaard diagrams and obtain a presentation for the fundamental group of the knot complement. In Section 3, we consider strongly-cyclic branched coverings of knots and give conditions for their

---

1991 *Mathematics Subject Classification.* Primary 57M12, 57M25; Secondary 20F05, 57M05, 20F10.

Key words and phrases. $(g,1)$-knots, $(g,1)$-decompositions, cyclic branched coverings, presentations of groups, Heegaard splittings, Heegaard diagrams.

Work performed under the auspices of G.N.S.A.G.A. of C.N.R. of Italy and supported by M.U.R.S.T., by the University of Bologna, funds for selected research topics, and by the Russian Foundation for the Basic Researches. The third named author was supported by the INTAS project "CalcoMet-GT' 03-51-3663 and by RFRB.

existence and uniqueness, using $(g,1)$-decompositions. As a relevant example we consider, in Section 4, the case of generalized periodic Takahashi manifolds. Section 5 is devoted to fundamental groups of strongly-cyclic branched coverings of knots. These groups admit geometric g-word cyclic presentations, a straightforward generalization of cyclic presentations of groups. Furthermore, we describe an algorithm to obtain such a presentation, starting from the group of the knot complement.

For the theory of Heegaard splittings of 3-manifolds we refer to [5].

2. $(g,1)$-DECOMPOSITIONS OF KNOTS

The notion of (g,b)-decomposition of a link L in a closed orientable 3-manifold has been introduced in [4], as a generalization of the classical bridge (or plat) decomposition of links in $\mathbf{S}^3$.

A non-empty set $\{a_1,\dots,a_b\}$ of mutually disjoint arcs properly embedded in a genus g handlebody $\mathcal{H}_g$ is called *trivial* if there exist b mutually disjoint discs $D_1,\dots,D_b \subset \mathcal{H}_g$ such that $a_i \cap D_i = a_i \cap \partial D_i = a_i$, $a_i \cap D_j = \emptyset$ and $\partial D_i - a_i \subset \partial\mathcal{H}_g$ for all $i,j=1,\dots,b$ and $i \neq j$.

Let $\mathcal{H}_g$ and $\mathcal{H}'_g$ be the two handlebodies of a genus g Heegaard splitting of a closed orientable 3-manifold N and let $S_g = \partial\mathcal{H}_g = \partial\mathcal{H}'_g$. A link $L \subset N$ is said to be in *b-bridge position* with respect to S_g if: (i) L intersects S_g transversally and (ii) $L \cap \mathcal{H}_g$ and $L \cap \mathcal{H}'_g$ are both sets of b mutually disjoint properly embedded trivial arcs. This splitting is called a *(g,b)-decomposition of L*.

Equivalently, a (g,b)-decomposition of L can be obtained by taking two genus g handlebodies $\mathcal{H}_g, \mathcal{H}'_g$, two trivial sets of arcs $A_b \subset \mathcal{H}_g$, $A'_b \subset \mathcal{H}'_g$ and an attaching homeomorphism $\varphi : (\partial\mathcal{H}'_g, \partial A'_b) \to (\partial\mathcal{H}_g, \partial A_b)$ such that

$$(N,L) \cong (\mathcal{H}_g, A_b) \cup_\varphi (\mathcal{H}'_g, A'_b).$$

Note that a $(0,b)$-decomposition is the classical bridge decomposition for links in a 3-sphere.

A link L is called a *(g,b)-link* if it admits a (g,b)-decomposition. Of course, a $(g,1)$-link is a knot, for every $g \geq 0$, and a $(0,1)$-knot is a trivial knot in $\mathbf{S}^3$. Obviously, a (g,b)-link is also a (g',b')-link for any $g' \geq g$ and $b' \geq b$.

In the following we shall restrict our attention to $(g,1)$-knots. A $(g,1)$-decomposition of a knot is represented in Figure 1.

A knot $K \subset N$ is called a *$(g,0)$-knot* if it can be embedded in a genus g Heegaard surface of N. It is easy to see that a $(g,0)$-knot is also a $(g,1)$-knot (just push an arc of K into a handlebody of the splitting and



FIGURE 1. A $(g,1)$-decomposition

the remaining part of the knot into the other handlebody). Moreover, when $b>0$, a (g,b)-knot is also a $(g+1,b-1)$-knot.

Remark 1. Let K be a knot in a 3-manifold N of Heegaard genus h. It is easy to see that, for any integer $b\geq 0$, there exists an integer $g\geq h$ such that K is a (g,b)-knot in N. Analogously, for any integer $g\geq h$ there exists an integer $b\geq 0$ such that K is a (g,b)-knot in N.

The following presentation theorem shows that a $(g,1)$-knot can be described by the images of a certain set of meridian curves.

Theorem 1. *Let $K\subset N$ be a knot with a $(g,1)$-decomposition $(N,K)=(\mathcal{H}_g,A)\cup_\varphi(\mathcal{H}'_g,A')$. Then:*

(i) *the knot K is completely determined, up to equivalence*[1]*, by $\varphi(\beta'_1),\varphi(\beta'_2),\dots,\varphi(\beta'_g)$, where $\beta'_1=\partial E'_1,\ \dots\ ,\ \beta'_g=\partial E'_g$, and $E'_1,\dots,E'_g$ is a complete system of meridian disks of $\mathcal{H}'_g$ not intersecting A';*

(ii) *if $(N,\bar{K})=(\mathcal{H}_g,A)\cup_{\bar{\varphi}}(\mathcal{H}'_g,A')$ is a $(g,1)$-decomposition of a knot $\bar{K}$ such that $\bar{\varphi}(\beta'_j)$ is isotopic to $\varphi(\beta'_j)$ in $\partial\mathcal{H}_g-\partial A$, for $j=1,\dots,g$, then $\bar{K}$ is equivalent to K;*

(iii) *if $A''\subset\partial\mathcal{H}_g$ is an arc such that $\partial A''=\partial A$ and $A''\cap\varphi(\beta'_j)=\emptyset$ for $j=1,\dots,g$, then $A\cup A''$ is a $(g,1)$-knot equivalent to K.*

Proof. (i) It follows from the fact that two properly embedded trivial arcs in a ball B, with the same endpoints, are isotopic rel ∂B. Statements (ii) and (iii) are straightforward. □

Now we find a presentation for the fundamental group of $(g,1)$-knots complements.

Let K be a knot in a manifold N with a $(g,1)$-decomposition $(N,K)=(\mathcal{H}_g,A)\cup_\varphi(\mathcal{H}'_g,A')$. Moreover, let $\tau:(\mathcal{H}_g,A)\to(\mathcal{H}'_g,A')$ be a fixed homeomorphism.

Consider the loops $\alpha_1,\dots,\alpha_g,\beta_1,\dots,\beta_g,\gamma$ in $\partial\mathcal{H}_g$, as depicted in Figure 2. Fix a base point $*$ on the curve γ and, for each $i=1,\dots,g$,

---

[1] Two knots $K\subset N$ and $K'\subset N'$ are considered *equivalent* if there exists a homeomorphism $\phi:N\to N'$ such that $\phi(K)=K'$.



FIGURE 2.

define the loops $\bar{\alpha}_i = \xi_i \cdot \alpha_i \cdot \xi_i^{-1}$ and $\bar{\beta}_i = \eta_i \cdot \beta_i \cdot \eta_i^{-1}$, where ξ_i and η_i are paths connecting $*$ to α_i and β_i, respectively. Obviously, the cycles $\bar{\alpha}_i$ and $\bar{\beta}_i$ are homologous to α_i and β_i, respectively, for each $i = 1, \ldots, g$. Moreover, we set $\bar{\alpha}'_i = \tau(\bar{\alpha}_i)$, $\bar{\beta}'_i = \tau(\bar{\beta}_i)$, for $i = 1, \ldots, g$, $\gamma' = \tau(\gamma)$ and $*' = \tau(*)$. Then $*' = \varphi^{-1}(*)$, up to isotopy. The homotopy classes of $\bar{\alpha}_1, \ldots, \bar{\alpha}_g, \bar{\beta}_1, \ldots, \bar{\beta}_g$ and γ generate $\pi_1(\partial\mathcal{H}_g - \partial A, *)$ and the homotopy classes of $\bar{\alpha}'_1, \ldots, \bar{\alpha}'_g, \bar{\beta}'_1, \ldots, \bar{\beta}'_g$ and γ' generate $\pi_1(\partial\mathcal{H}'_g - \partial A', *')$. In order to simplify the notation, in the following we use the same symbol for loops (resp. for cycles) and for corresponding homotopy classes (resp. homology classes).

Proposition 2. *The fundamental group of the complement of a knot* $K \subset N$*, with* $(g,1)$*-decomposition* $(N, K) = (\mathcal{H}_g, A) \cup_\varphi (\mathcal{H}'_g, A')$ *admits the presentation*

$$\pi_1(N - K, *) = \langle \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma \,|\, r_i(\bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma), \quad i = 1, \ldots, g\rangle, \tag{1}$$

where $r_i(\bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma) = \iota_\#\varphi_\#(\bar{\beta}'_i)$*, being* $\varphi_\#$ *and* $\iota_\#$ *the homomorphisms of fundamental groups induced by the attaching homeomorphism* $\varphi : (\partial\mathcal{H}'_g, \partial A') \to (\partial\mathcal{H}_g, \partial A)$ *and the inclusion* $\iota : (\partial\mathcal{H}_g, \partial A) \to (\mathcal{H}_g, A)$*, respectively.*

Proof. We have $\pi_1(\mathcal{H}_g - A, *) = \langle \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \bar{\beta}_1, \ldots, \bar{\beta}_g, \gamma \,|\, \bar{\beta}_1, \ldots, \bar{\beta}_g\rangle$ and $\pi_1(\mathcal{H}'_g - A', *') = \langle \bar{\alpha}'_1, \ldots, \bar{\alpha}'_g, \bar{\beta}'_1, \ldots, \bar{\beta}'_g, \gamma' \,|\, \bar{\beta}'_1, \ldots, \bar{\beta}'_g\rangle$. Applying the Seifert-van Kampen theorem to the $(g,1)$-decomposition of K, we get

$$\begin{aligned} \pi_1(N - K, *) = \langle & \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \bar{\beta}_1, \ldots, \bar{\beta}_g, \gamma, \bar{\alpha}'_1, \ldots, \bar{\alpha}'_g, \bar{\beta}'_1, \ldots, \bar{\beta}'_g, \gamma' \,|\\ & \bar{\beta}_i, \bar{\beta}'_i, \varphi(\bar{\alpha}'_i)\bar{\alpha}_i'^{-1}, \varphi(\bar{\beta}'_i)\bar{\beta}_i'^{-1}, \varphi(\gamma')\gamma'^{-1}, \quad i = 1, \ldots, g\rangle \\ = \langle & \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \bar{\beta}_1, \ldots, \bar{\beta}_g, \gamma \,|\, \bar{\beta}_i, \varphi(\bar{\beta}'_i), \quad i = 1, \ldots, g\rangle \\ = \langle & \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma \,|\, r_i(\bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma), \quad i = 1, \ldots, g\rangle, \end{aligned}$$

where $r_i(\bar{\alpha}_1, \ldots, \bar{\alpha}_g, \gamma)$ is obtained by erasing all $\bar{\beta}_j$'s terms from $\varphi(\bar{\beta}'_i)$. Since $\pi_1(\partial\mathcal{H}_g - \partial A, *) = \langle \bar{\alpha}_1, \ldots, \bar{\alpha}_g, \bar{\beta}_1, \ldots, \bar{\beta}_g, \gamma \,|\, \emptyset\rangle$, the statement is proved. □

As a consequence of the above proposition we have the following:

Corollary 3. *The first homology group of $N-K$ admits the following presentation:*

$$H_1(N-K)=\langle \alpha_1,\ldots,\alpha_g,\gamma \,|\, a_{i1}\alpha_1+\ldots+a_{ig}\alpha_g+b_i\gamma,\ i=1,\ldots,g\rangle \tag{2}$$

where, for each $i,j=1,\ldots,g$, a_{ij} (resp. b_i) is the exponent sum of $\bar{\alpha}_j$ (resp. γ) in the word r_i of the presentation (1).

3. Strongly-cyclic branched coverings of knots

Let $K\subset N$ be a knot in a 3-manifold N and $n>1$. Then an n-fold cyclic covering $f:M\to N$ branched over K is called *strongly-cyclic* if the branching index of K is n. This means that the fiber $f^{-1}(x)$ of each point $x\in K$ contains a single point. In this case the homology class of a meridian loop μ around K is mapped by the associated monodromy $\omega_f: H_1(N-K)\to\mathbb{Z}_n$ to a generator of the cyclic group $\mathbb{Z}_n$, and up to equivalence we can always suppose $\omega_f(\mu)=1$. We recall that two n-fold cyclic coverings $f':M'\to N$ and $f'':M''\to N$, branched over $K\subset N$, with monodromies $\omega_{f'}$ and $\omega_{f''}$ respectively, are equivalent if and only if there exists $u\in\mathbb{Z}_n$, with $\gcd(u,n)=1$, such that $\omega_{f''}=u\omega_{f'}$, where $u\omega_{f'}$ is the multiplication of $\omega_{f'}$ by u. For knots in $\mathbf{S}^3$, strongly-cyclic branched coverings and cyclic branched coverings are equivalent notions. The same property occurs in the general case when n is prime.

Now we discuss strongly-cyclic branched coverings of knots via their $(g,1)$-decompositions. In particular, we give existence conditions and compute their number, up to equivalence.

Let K be a knot in a 3-manifold N, and let

$$H_1(N)=\mathbb{Z}^d\oplus\mathbb{Z}_{t_1}\oplus\mathbb{Z}_{t_2}\oplus\cdots\oplus\mathbb{Z}_{t_s}, \tag{3}$$

with $0\le d$ and $1<t_1\,|\,t_2\,|\,\cdots\,|\,t_s$, be the canonical decomposition of the first integer homology group of N.

Let us fix a $(g,1)$-decomposition for K. Referring to presentation (2), let $H\in\mathcal{M}_g(\mathbb{Z})$ be the matrix with coefficients $h_{ij}=a_{ij}$ and $H'\in\mathcal{M}_{g,g+1}(\mathbb{Z})$ the augmented matrix with coefficients $h'_{ij}=a_{ij}$, for $i,j=1,\ldots,g$ and $h'_{ij}=b_i$ for $j=g+1$ and $i=1,\ldots,g$. Let $e_1,\ldots,e_r$ (resp. $e'_1,\ldots,e'_{r'}$) be the invariant factors of H (resp. H'). Moreover, set $e_{r+1}=\ldots=e_g=0$ and $e'_{r'+1}=\ldots=e'_g=0$.

Observe that H is a matrix presentation for $H_1(N)$. Therefore $s\le r$ and we have: $d=g-r$, $e_i=1$, for $1\le i\le r-s$, and $e_i=t_{i+s-r}$, for $r-s<i\le r$. Moreover, $H_1(N)$ is finite if and only if $\det H\ne 0$. In this case $\#H_1(N)=|\det H|$.

With the above notations, we have the following result.

Theorem 4. *Let K be a $(g,1)$-knot in a 3-manifold N with first homology group given by (3). Then K admits n-fold strongly-cyclic branched coverings if and only if* $\gcd(e_i, n) = \gcd(e'_i, n)$*, for each* $i = 1, \ldots, g$*. In this case the number of non-equivalent n-fold strongly-cyclic branched coverings of K is* $C_{N,n} = n^d \gcd(t_1, n) \gcd(t_2, n) \cdots \gcd(t_s, n)$*, which does not depend on K.*

Proof. Since we suppose $\omega_f(\gamma) = 1$, an n-fold strongly-cyclic branched covering of K is determined by a solution of the linear system (in $\mathbb{Z}_n$) $H\mathbf{x} = -\mathbf{b}$, and different solutions define non-equivalent coverings. So the result comes from [1], where systems of linear congruences are investigated. $\square$

As a consequence, we state a uniqueness result.

Corollary 5. *A knot K in a 3-manifold N admits a unique n-fold strongly-cyclic branched covering, up to equivalence, if and only if $H_1(N)$ is finite and* $\gcd(\#H_1(N), n) = 1$*.*

Proof. Let H be the matrix associated to a $(g,1)$-decomposition of K, according to the previous notations. If $H_1(N)$ is finite and $\gcd(\#H_1(N), n) = 1$, then H admits inverse in $\mathbb{Z}_n$, since in this case $|\det H| = \#H_1(N) \neq 0$. Therefore, $\mathbf{x} = -H^{-1}\mathbf{b}$ defines the unique n-fold strongly-cyclic branched covering of K. On the other hand, if K admits a unique n-fold strongly-cyclic branched covering, then $\gcd(e_i, n) = 1$ for $i = 1, \ldots, g$. Therefore $e_i \neq 0$, for each $i = 1, \ldots, g$. As a consequence, $|\det H| = e_1 e_2 \cdots e_g \neq 0$ and $\gcd(|\det H|, n) = 1$. Since in this case $\#H_1(N) = |\det H|$, the statement is proved. $\square$

The previous corollary holds, for any n, for knots in homology spheres (possibly $\mathbf{S}^3$), since in this case the first homology group is trivial.

Examples. We denote by M_k the connected sum of k copies of $\mathbf{S}^2 \times \mathbf{S}^1$, then $H_1(M_k) = \mathbb{Z}^k$.

- The trivial knot T in M_g is a $(g,1)$-knot defined by $\varphi(\beta'_i) = \beta_i$, for each $i = 1, \ldots, g$. So H and H' are both null matrices, and T admits exactly n^g non-equivalent n-fold strongly-cyclic branched coverings, each of them yielding M_{ng}.
- The "core" knot C in M_g is defined by $\varphi(\beta'_1) = \gamma$, and $\varphi(\beta'_i) = \beta_i$, for each $i = 2, \ldots, g$. In this case H is the null matrix but H' has $e'_1 = 1$, and therefore C admits no n-fold strongly-cyclic branched covering, for any $n > 1$.

4. The case of generalized periodic Takahashi manifolds

Generalized periodic Takahashi manifolds were constructed in [8] by Dehn surgery on a n-periodic $2mn$-component link, as illustrated in Figure 3, and denoted by $T_{n,m}(p_1/q_1, \dots, p_m/q_m; r_1/s_1, \dots, r_m/s_m)$, with $m, n > 0$, $p_i, q_i, r_i, s_i \in \mathbb{Z}$, $\gcd(p_i, q_i) = \gcd(r_i, s_i) = 1$, for each $i = 1, \dots, m$. For $m = 1$, we obtain the periodic Takahashi manifolds $T_n(p/q, r/s)$ introduced in [9].



Figure 3. $T_{n,m}(p_1/q_1, \dots, p_m/q_m; r_1/s_1, \dots, r_m/s_m)$

In [8] it is proved that generalized periodic Takahashi manifolds are strongly-cyclic branched coverings of certain knots in connected sums of lens spaces. That result is improved by the following.

Proposition 6. *For each* $n > 1$*, the generalized periodic Takahashi manifold* $T_{n,m}(p_1/q_1, \dots, p_m/q_m; r_1/s_1, \dots, r_m/s_m)$ *is an* n*-fold strongly-cyclic branched covering of a* $(2m, 0)$*-knot (and therefore of a* $(2m, 1)$*-knot).*

Proof. By [8, Theorem 8], $T_{n,m}(p_1/q_1, \dots, p_m/q_m; r_1/s_1, \dots, r_m/s_m)$ is an n-fold strongly-cyclic covering of the connected sum of lens spaces $N = L(p_1, q_1)\#L(r_1, s_1)\# \cdots \#L(p_m, q_m)\#L(r_m, s_m)$, obtained by rational surgery on a $2m$-component trivial link L, branched over a knot $K \subset N$, as shown in Figure 4.

Take $2m - 1$ "vertical" arcs $\sigma_1, \cdots, \sigma_{2m-1}$ consecutively connecting the components of L and consider a regular neighborhood $\mathcal{H}$ of $L \cup \sigma_1 \cup \cdots \cup \sigma_{2m-1}$. It is easy to see that $\partial\mathcal{H}$ is a genus $2m$ Heegaard surface of N, and K can be embedded on it, as depicted in Figure 5. Therefore K is $(2m, 0)$-knot and hence, as we already noted, a $(2m, 1)$-knot. □



Figure 4.

Now we explicitly give Heegaard diagrams of periodic Takahashi manifolds $T_n(p/q, r/s)$, for $p/q, r/s \geq 0$ (possibly equal to $+\infty = 1/0$). The cases $p/q < 0$ and/or $r/s < 0$ can be obtained in a similar way.

In Figure 6 an arc labelled k denotes k parallel arcs. Furthermore, each diagram in the figure admits a cyclic symmetry of order n; for $i = 1, \ldots, n$, we identify C_i (resp. D_i) with C'_i (resp. D'_i) by gluing together the first vertices of the marked ones, according to the orientations.

Proposition 7. *The periodic Takahashi manifold* $T_n(p/q, r/s)$*, for* $p/q, r/s \geq 0$ *(possibly equal to* $+\infty = 1/0$*), has Heegaard diagram as presented in Figure 6.*

Proof. Without loss of generality, we can assume that $p, q, r, s \geq 0$. Let $\mathcal{H}_2$ be a genus two handlebody canonically embedded in $\mathbb{R}^3$, as in Figure 7, which arises from Figure 5 for $m = 1$. Let us consider the Heegaard diagram of genus two of $N = L(p, q)\#L(r, s)$ obtained by cutting the genus two surface $\partial\mathcal{H}_2$ along the curves α_1 and α_2 and gluing them back after a twist of r/s and p/q turns, respectively, when $s, q \neq 0$. After this operation, the s (resp. q) "parallel" thin curves in $\partial\mathcal{H}_2$ become the curve $\varphi(\beta'_1)$ (resp. $\varphi(\beta'_2)$) of Theorem 1. The arcs $A'' \subset \partial\mathcal{H}_2$ and $A \subset \mathcal{H}_2$ (depicted in the figure by dashed lines) defines a knot $A \cup A''$ which is exactly the knot K of Proposition 6 for $m = 1$. If $s = 0$ (resp. $q = 0$) the curve $\varphi(\beta'_1)$ (resp. $\varphi(\beta'_2)$) is α_1 (resp. α_2).

By cutting $\mathcal{H}_2$ along β_1 and β_2 we obtain an open Heegaard diagram of genus two of N. From Corollary 3 we have $H_1(N - K) = \langle \alpha_1, \alpha_2, \gamma \,|\, p\alpha_2, r\alpha_1 \rangle$. So we have $z_1 = \omega_f(\alpha_1) = 0$ and $z_2 = \omega_f(\alpha_2) = 0$



FIGURE 5.

for each $n > 1$. By performing the n-fold strongly-cyclic covering determined by this monodromy, we obtain the Heegaard diagrams of Figure 6. □



Figure 6. Heegaard diagrams for $T_n(p/q, r/s)$



Figure 7.

5. Fundamental groups of strongly-cyclic branched coverings

It is shown in [7] that strongly-cyclic branched coverings of $(1,1)$-knots admit geometric (i.e. induced by Heegaard diagrams) cyclic presentations. In order to obtain a similar result for any knot, we give the following definition.

Definition. A balanced mn-generator presentation of a group $\langle X \mid R\,\rangle$ with generator set $X = \{x_{i,j} \mid 1 \le i \le m\,,\, 1 \le j \le n\}$ and relator set $R = \{r_{i,j} \mid 1 \le i \le m\,,\, 1 \le j \le n\}$ is said to be an *m-word cyclic presentation* if there exist m words $w_1, \ldots, w_m$ in the free group F_{mn} generated by X, such that the defining relations are of the form $r_{i,j} = \theta_n^{j-1}(w_i)$, with $j = 1, \ldots, n$, for each $i = 1, \ldots, m$, where $\theta_n : F_{mn} \to F_{mn}$ is the automorphism defined by $\theta_n(x_{i,j}) = x_{i,j+1}$ (with second subscript mod n), for each $j = 1, \ldots, n$ and $i = 1, \ldots, m$.

We denote this group by $G_n(w_1, \ldots, w_m)$. For $m = 1$ we have the classical case of cyclically presented group (see [6]).

There is a strict connection between strongly-cyclic branched coverings of $(g,1)$-knots and g-word cyclic presentations, as stated by the following:

Theorem 8. *Every n-fold strongly-cyclic branched covering M of a $(g,1)$-knot $K \subset N$ admits a Heegard splitting of genus gn with cyclic symmetry of order n, which induces a g-word cyclic presentation for the fundamental group of M.*

Proof. Since K is a $(g,1)$-knot in N, we have $(N,K) = (\mathcal{H}_g, A) \cup_\varphi (\mathcal{H}'_g, A')$, where $\varphi : (\partial\mathcal{H}'_g, \partial A') \to (\partial\mathcal{H}_g, \partial A)$ is an attaching homeomorphism. Let $f : (M, f^{-1}(K)) \to (N,K)$ be an n-fold strongly-cyclic branched covering of K. Then $\mathcal{H}_{gn} = f^{-1}(\mathcal{H}_g)$ and $\mathcal{H}'_{gn} = f^{-1}(\mathcal{H}'_g)$ are both handlebodies of genus gn. Moreover, $f^{-1}(A)$ and $f^{-1}(A')$ are both properly embedded arcs in $\mathcal{H}_{gn}$ and $\mathcal{H}'_{gn}$ respectively. Thus, we get a genus gn Heegaard splitting $(M, f^{-1}(K)) = (\mathcal{H}_{gn}, f^{-1}(A)) \cup_\Phi (\mathcal{H}'_{gn}, f^{-1}(A'))$, where $\Phi : \partial\mathcal{H}'_{gn} \to \partial\mathcal{H}_{gn}$ is the lift of φ with respect to f. The group $\pi_1(\mathcal{H}_g, *)$ is generated by the set $\{\bar{\alpha}_1, \ldots, \bar{\alpha}_g\}$ (see the notations introduced before Proposition 2). Let ω_f be the monodromy associated to the covering f, we can suppose that $\omega_f(\gamma) = 1$. Moreover, we can assume that $\omega_f(\bar{\alpha}_i) = 0$ for each $i = 1, \ldots, g$. In fact, if $\omega_f(\bar{\alpha}_i) = k_i$ we replace $\bar{\alpha}_i$ with a loop homotopic to $\bar{\alpha}_i\gamma^{-k_i}$. Taking a point $Q \in A$, let δ be an arc from $*$ to Q such that $\delta \cap A = Q$. Then $f^{-1}(Q)$ is a single point $\tilde{Q} \in f^{-1}(A)$ and $f^{-1}(*)$ consists of n points $\tilde{*}_1, \ldots, \tilde{*}_n$. For $j = 1, \ldots, n$ and $i = 1, \ldots, g$, let $\tilde{\alpha}_{ij}$ and $\tilde{\delta}_j$ be the lifts (with respect to f) of $\bar{\alpha}_i$ and δ respectively, both containing $\tilde{*}_j$. Then, for each $i = 1, \ldots, g$, the n loops

$$\alpha_{i1} = \tilde{\delta}_1^{-1} \cdot \tilde{\alpha}_{i1} \cdot \tilde{\delta}_1, \ \ldots, \ \alpha_{in} = \tilde{\delta}_n^{-1} \cdot \tilde{\alpha}_{in} \cdot \tilde{\delta}_n$$

are cyclically permutated by a generator Ψ of the group of covering transformations and the set $\{\alpha_{ij} \mid 1 \leq i \leq g\,,\, 1 \leq j \leq n\}$ generates $\pi_1(\mathcal{H}_{gn}, \tilde{Q})$. Let $\{E'_1, \ldots, E'_g\}$ be a complete system of meridian disks for $\mathcal{H}'_g$ such that $E'_i \cap A' = \emptyset$, for each $i = 1, \ldots, g$. Then $f^{-1}(E'_i)$ consists of n disks $\tilde{E}'_{i1}, \ldots, \tilde{E}'_{in}$, which are cyclically permutated by Ψ, for each $i = 1, \ldots, g$, and $f^{-1}(E'_1 \cup \cdots \cup E'_g)$ is a complete system of meridian disks for the handlebody $\mathcal{H}'_{gn}$. The curves $\Phi(\partial\tilde{E}'_{11}), \ldots, \Phi(\partial\tilde{E}'_{gn})$ give the relators for the presentation of $\pi_1(M, \tilde{Q})$ induced by the Heegaard splitting. Since, for each $i = 1, \ldots, g$, the generators $\alpha_{i1}, \ldots, \alpha_{in}$ and the relator curves $\partial\tilde{E}'_{i1}, \ldots, \partial\tilde{E}'_{in}$ are cyclically permutated by Ψ, each $\partial\tilde{E}'_{ij}$ corresponds to a cyclic permutation of the generators in the word w_i. □

Now we describe an algorithm to obtain a presentation of the fundamental group of a strongly-cyclic branched covering of a $(g, 1)$-knot.

Let ω_f be the monodromy of an n-fold strongly-cyclic branched covering of a $(g, 1)$-knot K in a 3-manifold N and let us consider the presentation of the knot group obtained in Proposition 2. Following the proof of Theorem 8, we choose, for each $i = 1, \ldots, g$, a new generator $\widehat{\alpha}_i = \bar{\alpha}_i\gamma^{-\omega_f(\bar{\alpha}_i)}$ and we get $\pi_1(N - K, *) = \langle \widehat{\alpha}_1, \ldots, \widehat{\alpha}_g, \gamma \,|\, \bar{r}_i(\widehat{\alpha}_1, \ldots, \widehat{\alpha}_g, \gamma), \quad i = 1, \ldots, g\rangle$, with $\bar{r}_i(\widehat{\alpha}_1, \ldots, \widehat{\alpha}_g, \gamma) = r_i(\widehat{\alpha}_1\gamma^{\omega_f(\bar{\alpha}_1)}, \ldots, \widehat{\alpha}_g\gamma^{\omega_f(\bar{\alpha}_g)}, \gamma)$.

Suppose now that, for each $i = 1, \dots, g$, we have

$$r_i(\widehat{\alpha}_1, \dots, \widehat{\alpha}_g, \gamma) = \widehat{\alpha}_{j_1}^{\varepsilon_{ij_1}} \gamma^{\eta_{ij_1}} \widehat{\alpha}_{j_2}^{\varepsilon_{ij_2}} \gamma^{\eta_{ij_2}} \dots \widehat{\alpha}_{j_{\lambda_i}}^{\varepsilon_{ij_{\lambda_i}}} \gamma^{\eta_{ij_{\lambda_i}}}$$

where $\varepsilon_{ij_l} \neq 0$ for each $l = 1, \dots, \lambda_i$ and $j_1, \dots, j_{\lambda_i}$ are not necessary distinct.

Then we have the following:

Proposition 9. *The fundamental group of the n-fold strongly-cyclic branched covering of a $(g,1)$-knot K, with monodromy ω_f, admits the g-word cyclic presentation $G_n(w_1, \dots, w_g)$, where, for each $i = 1, \dots, g$,*

$$w_i = x_{j_1,1}^{\varepsilon_{ij_1}} x_{j_2,k_{i2}}^{\varepsilon_{ij_2}} \cdots x_{j_{\lambda_i},k_{i\lambda_i}}^{\varepsilon_{ij_{\lambda_i}}},$$

with $k_{il} \equiv 1 + \sum_{h=1}^{l-1} \eta_{ij_h} \mod n$, for $l = 2, \dots, \lambda_i$.

Proof. Let M be the n-fold strongly-cyclic branched covering of K determined by ω_f. From the proof of Theorem 8, the fundamental group of M admits the set of generators $\{\alpha_{ij} \mid 1 \leq i \leq g\,,\, 1 \leq j \leq n\}$, where $\alpha_{i1}, \dots, \alpha_{in}$ are the components of the lifting of $\widehat{\alpha}_i$. The relators are (homotopic to) $\Phi(\partial\tilde{E}'_{11}), \dots, \Phi(\partial\tilde{E}'_{gn})$, where Φ is the lifting of φ with respect to f, and each $\partial\tilde{E}'_{ij}$ is a component of the lifting of a meridian disc E'_i of $\mathcal{H}'_g$ such that $E'_i \cap A' = \emptyset$. For each $i = 1, \dots, g$, we can choose E'_i such that $\partial E'_i = \tau(\beta_i)$, and therefore the relators are (homotopic to) the components of the lifting of $r_i(\bar{\alpha}_1, \dots, \bar{\alpha}_g, \gamma)$, or equivalently $\bar{r}_i(\widehat{\alpha}_1, \dots, \widehat{\alpha}_g, \gamma)$ $(i = 1, \dots, g)$, which arise from the relations $\bar{\beta}'_i = \varphi(\bar{\beta}'_i)$. Now, since $\omega_f(\gamma) = 1$, a factor γ^k lifts to a path connecting the point of $f^{-1}(*)$ in the sheet j with the corresponding point in the sheet $j + k$ (mod n). The result immediately follows. ☐

Example. Let us consider the $(2,1)$-knot $K = A \cup A''$ of Figure 7, with $q = s = 1$. In this case $K \subset N = L(p,1)\#L(r,1)$, and

$$\pi_1(N - K) = \langle \alpha_1, \alpha_2, \gamma \mid \alpha_2\alpha_1^{-r}\gamma\alpha_2^{-1}\gamma^{-1}, \alpha_1\gamma\alpha_2^{p}\alpha_1^{-1}\gamma^{-1} \rangle.$$

By applying Proposition 9, with $\omega_f(\alpha_1) = \omega_f(\alpha_2) = 0$, we get $\pi_1(T_n(p/1, r/1)) = G_n(w_1, w_2)$, where $w_1 = x_{2,1}x_{1,1}^{-r}x_{2,2}^{-1}$ and $w_2 = x_{1,1}x_{2,2}^{p}x_{1,2}^{-1}$, which is exactly the group presentation obtained in [9].

References

[1] A. T. Butson and B. M. Stewart: *Systems of linear congruences.* Canad. J. Math. **7** (1955), 358-368.

[2] A. Cattabriga and M. Mulazzani: *Strongly-cyclic branched coverings of* $(1,1)$-*knots and cyclic presentation of groups.* Math. Proc. Camb. Philos. Soc. **135** (2003), 137-146.

[3] A. Cattabriga and M. Mulazzani: $(1,1)$-*knots via the mapping class group of the twice punctured torus.* Adv. Geom. **4** (2004), 263-277.

[4] H. Doll. *A generalized bridge number for links in 3-manifold.* Math. Ann. **294** (1992), 701-717.

[5] J. Hempel. *3-manifolds.* Annals of Math. Studies, vol. 86, Princeton University Press, Princeton, N.J., 1976.

[6] D. L. Johnson. *Topics in the theory of group presentations.* London Math. Soc. Lect. Note Ser., vol. 42, Cambridge Univ. Press, Cambridge, U.K., 1980

[7] M. Mulazzani. *Cyclic presentations of groups and cyclic branched coverings of* $(1,1)$-*knots.* Bull. Korean Math. Soc. **40** (2003), 101-108.

[8] M. Mulazzani and A. Vesnin. *Generalized Takahashi manifolds.* Osaka J. Math. **39** (2002), 705-721.

[9] M. Takahashi. *On the presentations of the fundamental groups of 3-manifolds.* Tsukuba J. Math. **13** (1989), 175-189.

Department of Mathematics, University of Modena and Reggio Emilia, Italy
E-mail address: `cristofori.paola@unimo.it`

Department of Mathematics and C.I.R.A.M., University of Bologna, Italy
E-mail address: `mulazza@dm.unibo.it`

Sobolev Institute of Mathematics, Novosibirsk 630090, Russia
E-mail address: `vesnin@math.nsc.ru`